1. **Kobayashi H**, Shido Y: Combined supramalleolar osteotomy and ankle arthrodesis for end-stage ankle arthritis with large varus deformity. Foot Ankle Surg (in press).
2. **Kobayashi H**, Shido Y: Simultaneous hybrid closed-wedge supramalleolar osteotomy and ankle arthrodesis for posttraumatic varus ankle arthritis with periarticular deformity: A case report. Mod Rheumatol Case Rep 8:37-42, 2023.
3. **Kobayashi H**, Kageyama Y, Shido Y: Calcaneocuboid distraction arthrodesis with synthetic bone grafts: preliminary results of an innovative bone grafting procedure in 13 patients. J Foot Ankle Surg 56:1223-31, 2017.
4. **Kobayashi H**, Kageyama Y, Shido Y: Treatment of varus ankle osteoarthritis and instability with a novel mortise-plasty osteotomy procedure. J Foot Ankle Surg 55:60-7, 2016.
5. **Kobayashi H**, Kageyama Y, Shido Y: Gradual correction of traumatic hallux varus with metatarsal hemicallotasis. J Foot Ankle Surg 55:283-7, 2016.
6. **Kobayashi H**, Kageyama Y, Shido Y: Isolated fifth metatarsocuboid coalition: a case report. J Foot Ankle Surg 54:734-8, 2015.
7. Kageyama Y, Torikai E, Tsujimura K, **Kobayashi H**: Involvement of IL-33 in pathogenesis of rheumatoid arthritis: the effect of etanercept on the serum level of IL-33. Mod Rheumatol 22:89-93, 2012.
8. Ichikawa T, Kageyama Y, **Kobayashi H**, Kato N, Tsujimura K, Koide Y: Etanercept treatment reduces the serum levels of interleukin-15 and interferon-gamma inducible protein-10 in patients with rheumatoid arthritis. Rheumatol Int 30:725-30, 2010.
9. Kageyama Y, **Kobayashi H**, Kato N: Infliximab treatment reduces the serum levels of interleukin-23 in patients with rheumatoid arthritis. Mod Rheumatol 19:657-62, 2009.
10. Kageyama Y, **Kobayashi H**, Kato N, Shimazu M: Etanercept reduces the serum levels of macrophage chemotactic protein-1 in patients with rheumatoid arthritis. Mod Rheumatol 19:372-8, 2009.
11. Kageyama Y, Takahashi M, Nagafusa T, **Kobayashi H**, Nagano A: Reduction of urinary levels of pyridinoline and deoxypyridinoline and serum levels of soluble receptor activator of NF-kappa B ligand by etanercept in patients with rheumatoid arthritis. Clin Rheumatol 27:1093-101, 2008.
12. **Kobayashi H**, Sakurai M, Kobayashi T: Extensor digitorum longus tenosynovitis caused by talar head impingement in an ultramarathon runner. J Orthop Surg (Hong Kong) 15:245-7, 2007.
13. **Kobayashi H,** Hara J, Kitahama Y, Hanakita J: Osteoma arising from the lumbar articular process: a case report. Neurol Med Chir (Tokyo) 46:512-5, 2006.
14. **Kobayashi H**, Saito T, Koshino T: Immunolocalization of carboxy-terminal type Ⅱ procollagen peptide in regenerated articular cartilage of osteoarthritic knees after reduction of mechanical stress. Osteoarthritis Cartilage 10:870-8, 2002.